

Complete genome sequence of the motile actinomycete *Actinoplanes missouriensis* 431^T (= NBRC 102363^T)

Hideki Yamamura¹, Yasuo Ohnishi², Jun Ishikawa³, Natsuko Ichikawa⁴, Haruo Ikeda⁵, Mitsuo Sekine⁴, Takeshi Harada⁴, Sueharu Horinouchi^{2†}, Misa Otoguro^{1,6}, Tomohiko Tamura⁶, Ken-ichiro Suzuki⁶, Yasutaka Hoshino³, Akira Arisawa⁷, Youji Nakagawa¹, Nobuyuki Fujita⁴ and Masayuki Hayakawa¹

¹Division of Applied Biological Sciences, Interdisciplinary Graduate School of Medicine and Engineering, University of Yamanashi, Yamanashi, Japan

²Department of Biotechnology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan

³Department of Bioactive Molecules, National Institute of Infectious Diseases, Tokyo, Japan

⁴Biological Resource Center, National Institute of Technology and Evaluation, Tokyo, Japan

⁵Kitasato Institute for Life Sciences, Kitasato University, Kanagawa, Japan

⁶Biological Resource Center, National Institute of Technology and Evaluation, Chiba, Japan

⁷Bioresource Laboratories, MicroBiopharm Japan Co., Ltd., Shizuoka, Japan

Corresponding author: Masayuki Hayakawa (mhayakawa@yamanashi.ac.jp)

†Deceased July 12th, 2009

Actinoplanes missouriensis Couch 1963 is a well-characterized member of the genus *Actinoplanes*, which is of morphological interest because its members typically produce sporangia containing motile spores. The sporangiospores are motile by means of flagella and exhibit chemotactic properties. It is of further interest that members of *Actinoplanes* are prolific sources of novel antibiotics, enzymes, and other bioactive compounds. Here, we describe the features of *A. missouriensis* 431^T, together with the complete genome sequence and annotation. The 8,773,466 bp genome contains 8,125 protein-coding and 79 RNA genes.

Keywords: motile actinomycetes, sporangia, zoospores, motile spores, flagellation, aerobic, Gram-positive, *Micromonosporaceae*, *Actinoplanes*, *A. missouriensis*

Introduction

Strain 431^T [= NBRC 102363^T = DSM 43046^T = ATCC 14538^T and other culture collections] is the type strain of the species *Actinoplanes missouriensis* [1,2], which is a well-characterized member of the genus *Actinoplanes* [3]. The genus is of morphological interest because its members typically produce spherical, subspherical, cylindrical, or very irregular sporangia arising from vegetative mycelia [1,4]. Sporangiospores, which are released following the immersion of sporangia in water, are motile by means of polar or peritrichous flagella [5] and display a positive chemotactic response to a number of amino acids, aromatic compounds, sugars and inorganic ions [6,7]. Recently, the flagellin gene of various *Actinoplanes* strains was successfully amplified and sequenced to reveal the evolutionary relationships between flagellar genes [8]. As members

of the genus *Actinoplanes*, including *A. missouriensis*, produce a variety of antibiotics, enzymes and other bioactive compounds [9-12], the genomes of *Actinoplanes* species are potentially useful genetic resources for discovering secondary metabolites and enzymes. The genome analysis of the acarbose producer *Actinoplanes* sp. SE50/110 was most recently reported [13].

Here, we present a summary classification and a set of features for *A. missouriensis* strain 431^T, together with a description of the complete genome sequencing and annotation.

Classification and features

A. missouriensis strain 431^T was originally isolated from barnyard soil near Hamilton, Missouri, USA using the pollen-baiting technique and was first

described by Couch in 1963 [1]. Couch also isolated additional strains of *A. missouriensis* from ten soil collections obtained from the Mississippi Valley to the West Coast. The range of 16S rRNA gene sequence similarities between strain 431^T (= NBRC 102363^T) (AB711914) and valid members of the genus *Actinoplanes* was 95.7–97.9%. The highest sequence similarities were to *Actinoplanes utahensis* NBRC 13244^T.

Figure 1 shows the phylogenetic neighborhood of *A. missouriensis* 431^T in a 16S rRNA gene-based tree. The sequences of five of the six 16S rRNA gene copies in the genome of *A. missouriensis* 431^T

are identical (one sequence differed by 4 nucleotides), but differ by 55 nucleotides (3.8%) from the previously published 16S rRNA gene sequence of NBRC 13243^T (AB037008). The differences between the genome data and the reported 16S rRNA gene sequence are likely due to sequencing errors in the previously reported sequence data. The 16S rRNA sequence of strain 431^T did not match significantly with any 16S rRNA gene sequences from environmental genomic samples and surveys available at the NCBI BLAST server (March 2012).




Figure 1. Phylogenetic tree highlighting the position of *A. missouriensis* 431^T relative to *Actinoplanes* sp. strain SE50/110 and other type strains within the genus *Actinoplanes*. The tree was inferred from 1,351 aligned characters [14,15] of the 16S rRNA gene sequence under the maximum likelihood criterion [16] and rooted with the type strain of the neighboring genus *Dactylosporangium*. Only bootstrap values above 50% are shown (1,000 resamplings) at branching points. Lineages with type strain genome sequencing projects registered in GOLD [17] are labeled with an asterisk, those also listed as 'Complete and Published' with two asterisks.

The color of the substrate mycelia of strain 431^T is ochraceous salmon on Czapek agar. The strain does not form aerial mycelium. On Czapek agar, the strain produces a soluble pale lavender pigment [1]. Optimum growth occurs at 28°C. Strain 431^T produces globose spores (1-1.2 µm) arranged in irregular coils within a terminal sporangium. Sporangia are globose to subglobose in shape and 6 to 14 µm in diameter [Figure 2] [1].

Sporangia release motile spores, called zoospores, after unwrapping of the membranous sheath covering sporangia (Table 1) [5]. Released zoospores have chemotactic properties for several substrate types, including sugars, amino acids, aromatic compounds and mineral ions [6,7]. Flagella of the zoospores consist of a 44-kDa flagellar protein (FliC) [5] encoded by the *fliC* gene, which has been applied as a molecular marker in a taxonomic study of the genus *Actinoplanes* [8]. Strain 431^T utilizes D-xylose, L-arabinose, D-glucose, D-fructose, D-mannose, L-rhamnose, D-mannitol, sucrose, mannose, dextrin, D-galactose, D-lactose, methyl β-D-glucoside and L-rhamnose (1%, w/v),

but not *myo*-inositol, raffinose, cellulose, adonitol, D-arabinose, D-melezitose, methyl α-D-glucoside, D-ribose or D-sorbitol (1%, w/v) [4,29]. Strain 431^T also utilizes *p*-hydroxybenzoic acid, sodium acetate and sodium fumarate (0.1%, w/v), but not *m*-hydroxybenzoic acid, sodium succinate, syringic acid, or vanillin (0.1%, w/v) [4,29]. Strain 431^T degrades chitin, DNA, elastin, lecithin, RNA and tyrosine [26], and also actively decomposes natural rubber [32] and flavonoids, such as quercetin, rutin and hesperidin [33]. Strain 431^T produces glucose isomerase (xylose isomerase) with a molecular weight of approximately 8,000 daltons [34] and also produces 6-alkyl-4-*O*-dihydrogeranyl-2-methoxyhydroquinones as novel phenolic lipids [35]. Antimicrobial activity against *Streptomyces murinus* ISP 5091 is positive, but is negative against *Aspergillus niger* LIV 131, *Bacillus subtilis* NCIB 3610 and *Staphylococcus aureus* NCTC 8532 [29]. Additional physiological and drug susceptibility data are described in detail elsewhere [29].




Figure 2. Scanning electron micrograph of *A. missouriensis* 431^T. Bar interval, 10 µm.

Table 1. Classification and general features of *A. missouriensis* 431^T based on the MIGS recommendations [18]

MIGS ID	Property	Term	Evidence code
		Domain <i>Bacteria</i>	TAS [19]
		Phylum <i>Actinobacteria</i>	TAS [20]
		Class <i>Actinobacteria</i>	TAS [21]
	Current classification	Order <i>Actinomycetales</i>	TAS [2,21-24]
		Family <i>Micromonosporaceae</i>	TAS [2,21,24-26]
		Genus <i>Actinoplanes</i>	TAS [2,3,27,28]
		Species <i>Actinoplanes missouriensis</i>	TAS [1,2]
		Type strain 431	TAS [1,2]
	Gram stain	Not tested, likely positive	NAS
	Cell shape	Produces sporangium	TAS [1]
	Motility	Motile	TAS [1]
	Sporulation	Motile spore	TAS [1]
	Temperature range	Mesophile, temperature range not determined	TAS [29,30]
	Optimum temperature	28°C	TAS [29,30]
	Salinity	2% NaCl	TAS [29]
MIGS-22	Oxygen requirement	Aerobe	NAS
	Carbon source	Several (refer to text)	TAS [29]
	Energy source	Carbohydrates	TAS [29]
MIGS-6	Habitat	Soil	TAS [1]
MIGS-15	Biotic relationship	Free-living	NAS
MIGS-14	Pathogenicity	None	NAS
	Biosafety level	1	NAS
	Isolation	Barnyard soil	TAS [1]
MIGS-4	Geographic location	Hamilton, Missouri, USA	TAS [1]
MIGS-5	Sample collection time	1963 or before	TAS [1]
MIGS-4.1	Latitude	Not reported	
MIGS-4.2	Longitude	Not Reported	
MIGS-4.3	Depth	Not reported	
MIGS-4.4	Altitude	Not reported	

Evidence codes - IDA: Inferred from Direct Assay (first time in publication); TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [31]. If the evidence code is IDA, then the property was directly observed by one of the authors or an expert mentioned in the acknowledgements.

Chemotaxonomy

The diagnostic amino acids of strain 431^T are meso- and hydroxydiaminopimelic acids, and the characteristic whole-cell sugars are xylose and arabinose [29,36]. The peptidoglycan of strain 431^T contains glycosylated muramic acid residues [28,37], and the major phospholipids are phosphatidylethanolamine, diphosphatidylglycerol, phosphatidylglycerol, phosphatidylinositol and phosphatidylinositol mannoside [28]. Major fatty acids (relative ratio %) are iso-C_{16:0} (18.3) and C_{19:0} (26.4) [28,29]. The predominant menaquinone is MK-9(H₄) (>50%), and the minor menaquinones are MK-9(H₆) and MK-10(H₄) [28,29].

Genome sequencing information

Genome project history

A consortium consisting of universities, research institutions and private companies in Japan was organized to accelerate cooperative research and development utilizing genome sequences of actinomycetes. The consortium successfully accomplished the genome projects of *Kitasatospora setae* NBRC 14216^T [38] and *A. missouriensis* 431^T (= NBRC 102363^T), as reported here. The complete genome sequences of these strains have been deposited in the INSDC database and are also available from the DOGAN genome database [39]. A summary of the genome sequencing project information of *A. missouriensis* is shown in Table 2.

Growth conditions and DNA isolation

A. missouriensis strain 431^T (= NBRC 102363^T) was grown in NBRC 231 medium (Maltose-Bennett's Agar) at 28°C. DNA was isolated from 3.76 g of mycelial paste using the CTAB method and fragmented with the HydroShear device (Genomic Solutions), as recommended by the manufacturer.

Genome sequencing and assembly

The genome sequence of *A. missouriensis* 431^T was determined using a whole-genome shotgun sequencing approach together with the Sanger method. DNA shotgun libraries with average insert sizes of 1.6 and 6 kbp were constructed in pUC118, and a library with an average insert size of 36 kbp was constructed in pCC1FOS (Epicentre). A total of 96,384 reads were trimmed at a threshold quality value of 20 and assembled using the Phrap and CONSED assembly tools [40,41]. A total of 141 gaps (18,532 bp in total) were closed by sequencing PCR products, and 236 low-quality regions were re-sequenced by primer walking. In the final assembly step, we confirmed that each base of the genome was sequenced from multiple clones in either both or a single direction and had Phrap quality scores of >70 and >40, respectively. For validation of the contig alignment, the Argus Optical Mapping System (OpGen, Inc.) was used.

Table 2. Genome sequencing project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	Complete Three Sanger libraries: 1.5- and 6-kb pUC118, and 36-kb fosmid pCC1Fos
MIGS-28	Libraries used	
MIGS-29	Sequencing platform	Sanger ABI 3730xl
MIGS-31.2	Fold coverage	8.7 ×
MIGS-30	Assemblers	Phrap
MIGS-32	Gene calling method	Glimmer3, tRNAscan-SE 1.23
	Genome database release	DOGAN [39]
	Genbank ID	AP012319
	Genbank date of release	2012/03/31
	GOLD ID	Gc02182
	Project relevance	Biotechnological

Genome annotation

Gene prediction was performed using Glimmer3 [42] and tRNA-scanSE [43], followed by manual inspection of each translation start site using the Frameplot program [44]. Similarity search results against the NCBI nr and Pfam databases were used for functional prediction. Manual functional annotation was performed within an in-house platform developed by J.I. (Unpublished).

Genome properties

The genome of strain 431^T consists of one circular chromosome with a length of 8,773,466 bp and a G+C content of 70.8% (Table 3 and Figure 3). *A. missouriensis* DSM 43046^T was reported to contain a linear plasmid, pAM1 [33]; however, the corresponding plasmid was not found in strain 431^T (=

NBRC 102363^T) by CHEF electrophoresis analysis. It is possible that the linear plasmid was cured from strain 431^T by repeated subculturing in the laboratory. In comparison to other members of the family *Micromonosporaceae*, such as *Salinispora* and *Micromonospora*, *A. missouriensis* has a larger genome size. Of the total of 8,204 predicted genes, 8,125 were protein-coding genes and 79 were RNA genes. More than half of the protein-coding genes (4,539, 55.9%) were assigned a putative function, while the remaining predicted genes were annotated as hypothetical proteins. The distribution of genes into COGs functional categories is presented in Table 4.

Table 3. Genome statistics

Attribute	Value	% of total ^a
Genome size (bp)	8,773,466	100.00%
Coding region (bp)	7,964,669	90.78%
G+C content (bp)	6,213,639	70.82%
Total genes	8,204	n/a
RNA genes	79	n/a
rRNA operon	6	n/a
tRNA (44 species)	58	n/a
tmRNA	1	n/a
4.5S RNA	1	n/a
<i>rnpB</i> (RNase P component)	1	n/a
Protein-coding genes	8,125	100.00%
Genes in paralog clusters	3,123	38.44%
Genes assigned to COGs	5,171	63.64%
Genes assigned Pfam domains	5,463	67.24%
Genes with signal peptides ^b	799	9.83%
Genes with transmembrane helices	2,122	26.12%

a) The total is based on either the size of the genome in base pairs or the total number of protein-coding genes in the annotated genome.

b) Predicted by SignalP 4.0.

n/a, not available




Figure 3. Graphical circular map of the strain 431^T genome. From outside to the center: genes on forward strand (color by COG categories), Genes on reverse strand (color by COG categories), RNA genes (tRNA genes, green; rRNA genes, red; other RNA gene types, black), GC content, GC skew.

Table 4. Number of genes associated with the 25 general COG functional categories

Code	Value	%age^a	Description
J	199	2.4	Translation, ribosomal structure and biogenesis
A	1	0	RNA processing and modification
K	831	10.2	Transcription
L	234	2.9	Replication, recombination and repair
B	1	0	Chromatin structure and dynamics
D	42	0.5	Cell cycle control, mitosis and meiosis
Y	0	0	Nuclear structure
V	143	1.8	Defense mechanisms
T	528	6.5	Signal transduction mechanisms
M	240	3.0	Cell wall/membrane biogenesis
N	80	1.0	Cell motility
Z	1	0	Cytoskeleton
W	0	0	Extracellular structures
U	52	0.6	Intracellular trafficking and secretion
O	156	1.9	Posttranslational modification, protein turnover, chaperones
C	360	4.4	Energy production and conversion
G	579	7.1	Carbohydrate transport and metabolism
E	524	6.4	Amino acid transport and metabolism
F	91	1.1	Nucleotide transport and metabolism
H	199	2.4	Coenzyme transport and metabolism
I	286	3.5	Lipid transport and metabolism
P	326	4.0	Inorganic ion transport and metabolism
Q	236	2.9	Secondary metabolites biosynthesis, transport and catabolism
R	976	12.0	General function prediction only
S	387	4.8	Function unknown
-	2,954	36.4	Not in COGs

a) The total is based on the total number of protein-coding genes in the annotated genome.

References

1. Couch JN. Some new genera and species of the *Actinoplanaceae*. *J Elisha Mitchell Sci Soc* 1963; **79**:53-70.
2. Skerman VBD, McGowan V, Sneath PHA. Approved lists of bacterial names. *Int J Syst Bacteriol* 1980; **30**:225-420.
<http://dx.doi.org/10.1099/00207713-30-1-225>
3. Couch JN. *Actinoplanes*, a new genus of the *Actinomycetales*. *J Elisha Mitchell Sci Soc* 1950; **66**:87-92.
4. Vobis G. Actinoplanetes. In: ST Williams (eds) *Bergey's Manual of Systematic Bacteriology*, Volume 4, Williams & Winkins. Baltimore, 1989, p. 2418–2450.
5. Uchida K, Jang MS, Ohnishi Y, Horinouchi S, Hayakawa M, Fujita N, Aizawa S. Characterization of *Actinoplanes missouriensis* spore flagella. *Appl Environ Microbiol* 2011; **77**:2559-2562. PubMed <http://dx.doi.org/10.1128/AEM.02061-10>
6. Arora DK. Chemotaxis of *Actinoplanes missouriensis* zoospores to fungal conidia, chlamydospores and sclerotia. *J Gen Microbiol* 1986; **132**:1657-1663.
7. Hayakawa M, Tamura T, Nonomura H. Selective isolation of *Actinoplanes* and *Dactylosporangium* from soil by using γ -collidine as the

- chemoattractant. *J Ferment Bioeng* 1991; **72**:426-432. [http://dx.doi.org/10.1016/0922-338X\(91\)90049-M](http://dx.doi.org/10.1016/0922-338X(91)90049-M)
8. Yamamura H, Hanawa K, Kusunoki M, Nakamura K, Nakagawa Y, Ishida Y, Hamada M, Otoguro M, Tamura T, Fujita N, et al. Variation and predicted structure of the flagellin gene in *Actinoplanes* species. *FEMS Microbiol Lett* 2011; **324**:21-27. [PubMed](#) <http://dx.doi.org/10.1111/j.1574-6968.2011.02380.x>
 9. Amore R, Hollenberg CP. Xylose isomerase from *Actinoplanes missouriensis*: primary structure of the gene and protein. *Nucleic Acids Res* 1989; **17**:7515. [PubMed](#) <http://dx.doi.org/10.1093/nar/17.18.7515>
 10. Jarling M, Cauvet T, Grundmeire M, Kuhnert K, Pape H. Isolation of *mak1* from *Actinoplanes missouriensis* and evidence that Pep2 from *Streptomyces coelicolor* is a maltokinase. *J Basic Microbiol* 2004; **44**:360-373. [PubMed](#) <http://dx.doi.org/10.1002/jobm.200410403>
 11. Debono M, Merkel KE, Molloy RM, Barnhart M, Presti E, Hunt AH, Hamill RL. Actaplanin, new glycopeptide antibiotics produced by *Actinoplanes missouriensis*. The isolation and preliminary chemical characterization of actaplanin. *J Antibiot (Tokyo)* 1984; **37**:85-95. [PubMed](#) <http://dx.doi.org/10.7164/antibiotics.37.85>
 12. Ensign JC. Formation, properties, and germination of actinomycete spores. *Annu Rev Microbiol* 1978; **32**:185-219. [PubMed](#) <http://dx.doi.org/10.1146/annurev.mi.32.100178.001153>
 13. Schwientek P, Szczepanowski R, Rückert C, Kalinowski J, Klein A, Selber K, Wehmeier UF, Stoye J, Pühler A. The complete genome sequence of the acarbose producer *Actinoplanes* sp. SE50/110. *BMC Genomics* 2012; **13**:112. [PubMed](#) <http://dx.doi.org/10.1186/1471-2164-13-112>
 14. Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; **17**:540-552. [PubMed](#) <http://dx.doi.org/10.1093/oxfordjournals.molbev.a026334>
 15. Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics* 2002; **18**:452-464. [PubMed](#) <http://dx.doi.org/10.1093/bioinformatics/18.3.452>
 16. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011; **28**:2731-2739. [PubMed](#) <http://dx.doi.org/10.1093/molbev/msr121>
 17. Liolios K, Chen IM, Mavromatis K, Tavernarakis N, Hugenholtz P, Markowitz VM, Kyropides NC. The Genomes On Line Database (GOLD) in 2009: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res* 2010; **38**:D346-D354. [PubMed](#) <http://dx.doi.org/10.1093/nar/gkp848>
 18. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](#) <http://dx.doi.org/10.1038/nbt1360>
 19. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](#) <http://dx.doi.org/10.1073/pnas.87.12.4576>
 20. Garrity GM, Holt JG. The Road Map to the Manual. In: Garrity GM, Boone DR, Castenholz RW (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Springer, New York, 2001, p. 119-169.
 21. Stackebrandt E, Rainey F, Ward-Rainey N. Proposal for a new hierachic classification system, *Actinobacteria* classis nov. *Int J Syst Bacteriol* 1997; **47**:479-491. <http://dx.doi.org/10.1099/00207713-47-2-479>
 22. Buchanan RE. Studies in the Nomenclature and classification of the Bacteria: VIII. The subgroups and genera of the *Actinomycetales*. *J Bacteriol* 1918; **3**:403-406. [PubMed](#)
 23. Buchanan RE. Studies in the nomenclature and classification of bacteria. II. The primary subdivisions of the Schizomycetes. *J Bacteriol* 1917; **2**:155-164. [PubMed](#)
 24. Zhi XY, Li WJ, Stackebrandt E. An update of the structure and 16S rRNA gene sequence-based definition of higher ranks of the class *Actinobacteria*, with the proposal of two new suborders and four new families and emended descriptions of the existing higher taxa. *Int J Syst Evol Microbiol* 2009; **59**:589-608. [PubMed](#) <http://dx.doi.org/10.1099/ijss.0.65780-0>
 25. Krasil'nikov NA. Ray fungi and Related Organisms. *Actinomycetales*, Akademii Nauk, Moscow, 1938, p. 272.

26. Koch C, Kroppenstedt RM, Rainey FA, Stackebrandt E. 16S ribosomal DNA analysis of the genera *Micromonospora*, *Actinoplanes*, *Catellatospora*, *Catenuloplanes*, *Couchioplanes*, *Dactylosporangium*, and *Pilimelia* and emendation of the family *Micromonosporaceae*. *Int J Syst Bacteriol* 1996; **46**:765-768. [PubMed](#) <http://dx.doi.org/10.1099/00207713-46-3-765>
27. Couch JN, Bland CE. Genus I. *Actinoplanes* Couch 1950, 89; emend. mut. char. 1955, 153. In: Buchanan RE, Gibbons NE (eds), Bergey's Manual of Determinative Bacteriology, Eighth Edition, The Williams and Wilkins Co., Baltimore, 1974, p. 708-710.
28. Stackebrandt E, Kroppenstedt RM. Union of the genera *Actinoplanes* Couch, *Ampullariella* Couch, and *Amorphosporangium* Couch in a redefined genus *Actinoplanes*. *Syst Appl Microbiol* 1987; **9**:110-114. [http://dx.doi.org/10.1016/S0723-2020\(87\)80063-2](http://dx.doi.org/10.1016/S0723-2020(87)80063-2)
29. Goodfellow M, Stanton LJ, Simpson KE, Minnikin DE. Numerical and chemical classification of *Actinoplanes* and some related actinomycetes. *J Gen Microbiol* 1990; **136**:19-36. <http://dx.doi.org/10.1099/00221287-136-1-19>
30. Wink JM. Compendium of Actinobacteria. http://www.dsmz.de/microorganisms/wink_pdf/DSM43247.pdf
31. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene Ontology: tool for the unification of biology. *Nat Genet* 2000; **25**:25-29. [PubMed](#) <http://dx.doi.org/10.1038/75556>
32. Jendrossek D, Tomasi G, Kroppenstedt RM. Bacterial degradation of natural rubber: a privilege of actinomycetes? *FEMS Microbiol Lett* 1997; **150**:179-188. [PubMed](#) [http://dx.doi.org/10.1016/S0378-1097\(97\)00072-4](http://dx.doi.org/10.1016/S0378-1097(97)00072-4)
33. Rose K, Fetzner S. Identification of linear plasmid pAM1 in the flavonoid degrading strain *Actinoplanes missouriensis*^T (DSM 43046). *Plasmid* 2006; **55**:249-254. [PubMed](#) <http://dx.doi.org/10.1016/j.plasmid.2005.10.003>
34. Gong CS, Chen LF, Tsao GT. Purification and properties of glucose isomerase of *Actinoplanes missouriensis*. *Biotechnol Bioeng* 1980; **22**:833-845. <http://dx.doi.org/10.1002/bit.260220408>
35. Awakawa T, Fujita N, Hayakawa M, Ohnishi Y, Horinouchi S. Characterization of the biosynthesis gene cluster for alkyl-O-dihydrogeranyl-methoxyhydroquinones in *Actinoplanes missouriensis*. *ChemBioChem* 2011; **12**:439-448. [PubMed](#) <http://dx.doi.org/10.1002/cbic.201000628>
36. Horan AC, Brodsky B. *Actinoplanes caeruleus* sp. nov., a blue-pigmented species of the genus *Actinoplanes*. *Int J Syst Bacteriol* 1986; **36**:187-191. <http://dx.doi.org/10.1099/00207713-36-2-187>
37. Kawamoto I, Oka T, Nara T. Cell wall composition of *Micromonospora olivoasterospora*, *Micromonospora sagamiensis*, and related organisms. *J Bacteriol* 1981; **146**:527-534. [PubMed](#)
38. Ichikawa N, Oguchi A, Ikeda H, Ishikawa J, Kitani S, Watanabe Y, Nakamura S, Katano Y, Kishi E, Sasagawa M, et al. Genome sequence of *Kitasatospora setae* NBRC 14216^T: an evolutionary snapshot of the family *Streptomycetaceae*. *DNA Res* 2010; **17**:393-406. [PubMed](#) <http://dx.doi.org/10.1093/dnares/dsq026>
39. DOGAN genome database. <http://www.bio.nite.go.jp/dogan>.
40. Ewing B, Hillier L, Wendl MC, Green P. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res* 1998; **8**:175-185. [PubMed](#)
41. Ewing B, Green P. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res* 1998; **8**:186-194. [PubMed](#)
42. Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 2007; **23**:673-679. [PubMed](#) <http://dx.doi.org/10.1093/bioinformatics/btm009>
43. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 1997; **25**:955-964. [PubMed](#)
44. Ishikawa J, Hotta K. FramePlot: a new implementation of the frame analysis for predicting protein-coding regions in bacterial DNA with a high G + C content. *FEMS Microbiol Lett* 1999; **174**:251-253. [PubMed](#) <http://dx.doi.org/10.1111/j.1574-6968.1999.tb13576.x>