

EXTENDED GENOME REPORT

Open Access



High-quality draft genome sequence of *Ensifer meliloti* Mlalz-1, a microsymbiont of *Medicago laciniata* (L.) miller collected in Lanzarote, Canary Islands, Spain

Wan Adnawani Meor Osman¹, Peter van Berkum², Milagos León-Barrios³, Encarna Velázquez⁴, Patrick Elia², Rui Tian¹, Julie Ardley¹ , Margaret Gollagher⁵, Rekha Seshadri⁶, T. B. K. Reddy⁶, Natalia Ivanova⁶, Tanja Woyke⁶, Amrita Pati⁷, Victor Markowitz⁷, Mohamed N. Baeshen⁸, Naseebh Nabeeh Baeshen⁸, Nikos Kyrpides⁶ and Wayne Reeve^{1*}

Abstract

Ensifer meliloti Mlalz-1 (INSDC = ATZD00000000) is an aerobic, motile, Gram-negative, non-spore-forming rod that was isolated from an effective nitrogen-fixing nodule of *Medicago laciniata* (L.) Miller from a soil sample collected near the town of Guatiza on the island of Lanzarote, the Canary Islands, Spain. This strain nodulates and forms an effective symbiosis with the highly specific host *M. laciniata*. This rhizobial genome was sequenced as part of the DOE Joint Genome Institute 2010 Genomic Encyclopedia for Bacteria and Archaea-Root Nodule Bacteria (GEBA-RNB) sequencing project. Here the features of *E. meliloti* Mlalz-1 are described, together with high-quality permanent draft genome sequence information and annotation. The 6,664,116 bp high-quality draft genome is arranged in 99 scaffolds of 100 contigs, containing 6314 protein-coding genes and 74 RNA-only encoding genes. Strain Mlalz-1 is closely related to *Ensifer meliloti* IAM 12611^T, *Ensifer medicae* A 321^T and *Ensifer numidicus* ORS 1407^T, based on 16S rRNA gene sequences. gANI values of $\geq 98.1\%$ support the classification of strain Mlalz-1 as *E. meliloti*. Nodulation of *M. laciniata* requires a specific *nodC* allele, and the *nodC* gene of strain Mlalz-1 shares $\geq 98\%$ sequence identity with *nodC* of *M. laciniata*-nodulating *Ensifer* strains, but $\leq 93\%$ with *nodC* of *Ensifer* strains that nodulate other *Medicago* species. Strain Mlalz-1 is unique among sequenced *E. meliloti* strains in possessing genes encoding components of a T2SS and in having two versions of the adaptive acid tolerance response *lpiA-acvB* operon. In *E. medicae* strain WSM419, *lpiA* is essential for enhancing survival in lethal acid conditions. The second copy of the *lpiA-acvB* operon of strain Mlalz-1 has highest sequence identity ($> 96\%$) with that of *E. medicae* strains, which suggests genetic recombination between strain Mlalz-1 and *E. medicae* and the horizontal gene transfer of *lpiA-acvB*.

Keywords: Root-nodule bacteria, *Ensifer*, Geba-Rnb, *Medicago*, *lpiA-acvB* operon

Introduction

Symbiotic nitrogen fixation by pasture legumes and their associated root nodule bacteria provides a critical contribution to sustainable animal and plant production, and the maintenance of soil fertility in agricultural systems [1–3]. As such, it is of direct relevance to maintaining environmentally sustainable high agricultural yields,

which significantly contributes to the Sustainable Development Goals adopted in September 2015 as part of the UN's development agenda 'Transforming our world: the 2030 Agenda for Sustainable Development' [4]. Medics (*Medicago* spp.) are some of the most important and extensively grown pasture legumes and their specific symbiosis with strains of rhizobia belonging to either *Ensifer* (synonym *Sinorhizobium*) *meliloti* or the closely related species *E. medicae* [5, 6] has been the subject of extensive research efforts [7].

* Correspondence: W.Reeve@murdoch.edu.au

¹School of Veterinary and Life Sciences, Murdoch University, Murdoch, WA, Australia

Full list of author information is available at the end of the article



Medicago laciniata (L.) Miller (cut leaf medic), an annual native of southern and eastern Mediterranean and Saharo-Sindian countries, is of importance because of its ability to grow in comparatively arid habitats and marginal cropping areas [8–11]. It is highly specific in its rhizobial requirements, forming a symbiosis only with a restricted subset of *E. meliloti* and not with strains that nodulate *Medicago sativa* L. (alfalfa) or *Medicago truncatula* Gaertn. [12, 13]. This symbiotic specificity has been linked to the rhizobial *nod* genes, in particular a specific *nodC* allele [14]. For example, van Berkum and colleagues found that most rhizobial strains isolated from Tunisian *M. truncatula* and *M. laciniata* shared chromosomal identity, but differed in their *nodC* alleles [15]. Based on these and other differing symbiotic traits, Villegas et al. [13] proposed two biovars within *E. meliloti*: bv. medicaginis for *Ensifer* strains that are symbiotically efficient on *M. laciniata* and bv. meliloti for the classical *E. meliloti* group that efficiently nodulates *M. sativa*. However, in subsequent studies the diversity observed within bv. medicaginis strains indicate that this group is certainly heterogeneous [16].

M. laciniata is native to the Canary Islands and is present on all of the islands of this archipelago, growing in environments that range from arid to subhumid. *Ensifer meliloti* strain Mlalz-1 was isolated from a N₂-fixing nodule of *M. laciniata* grown in alkaline soil (pH 9.0) collected in Guatiza, in the arid Northeast of Lanzarote Island, in 2007. This strain was one of the rhizobial genomes sequenced as part of the DOE Joint Genome Institute 2010 GEBA-RNB project proposal [17, 18]. Here an analysis of the complete genome sequence of *E. meliloti* Mlalz-1 is provided.

Organism information

Classification and features

E. meliloti Mlalz-1 is a motile, non-sporulating, non-encapsulated, Gram-negative strain in the class *Alpha-proteobacteria*. The rod shaped form has dimensions of

approximately 0.5 μm in width and 1.0–2.0 μm in length (Fig. 1 *Left* and *Center*). It is fast growing, forming colonies after 3–5 days when grown on ½LA, TY, or a modified yeast-mannitol agar [19] at 28 °C. Colonies on ½LA are opaque, slightly domed and moderately mucoid with smooth margins (Fig. 1 *Right*). Minimum Information about the Genome Sequence (MIGS) for strain Mlalz-1 is provided in Table 1 and Additional file 1: Table S1.

Symbiotaxonomy

M. laciniata is a highly specific host and its microsymbionts also appear to be highly specific since studies of *Medicago* isolates have shown that *M. laciniata* strains fail to nodulate a range of *Medicago* species [5, 12]. Bailly et al. [20] reported that isolates of *M. laciniata* nodulated and fixed nitrogen with *M. truncatula*, but also provided evidence that these were the progeny of horizontal transfer of the nodulation genes. Strain Mlalz-1 nodulates and is effective for nitrogen fixation with *M. laciniata*. We report here that strain Mlalz-1 is unable to nodulate *Medicago polymorpha* L., the definitive host for *E. medicae* strains [6].

Extended feature descriptions

Previous studies using multilocus sequence typing showed that *M. laciniata* rhizobia did not form a distinct chromosomal group [15]. Phylogenetic analysis of strain Mlalz-1 was performed by aligning the 16S rRNA sequence (1389 bp from scaffold 84.85) to the 16S rRNA gene sequences of *Ensifer* type strains (Fig. 2). Based on four variable sites within this 16S rRNA gene sequence alignment, strain Mlalz-1 is closely related to *E. meliloti* IAM 12611^T (= LMG 6133^T) [21], *E. medicae* A 321^T (= LMG 19920^T) [6] and *E. numidicus* ORS 1407^T [22]. The available IMG 16S rRNA sequence of strain Mlalz-1 gave alignment identities of 100% to *E. meliloti* IAM 12611^T, 99.7% to *E. medicae* A 321^T and 99.5% to *E. numidicus* ORS 1407^T. In contrast, *E. meliloti* IAM




Fig. 1 Images of *Ensifer meliloti* Mlalz-1 using scanning (Left (a)) and transmission (Center (b)) electron microscopy as well as light microscopy to visualize colony morphology on solid media (Right (c))

Table 1 Classification and general features of *Ensifer meliloti* Mlalz-1 in accordance with the MIGS recommendations [65] published by the Genomic Standards Consortium [66]

MIGS ID	Property	Term	Evidence code ^a
	Current classification	Domain Bacteria	TAS [67]
		Phylum <i>Proteobacteria</i>	TAS [68]
		Class <i>Alphaproteobacteria</i>	TAS [69, 70]
		Order <i>Rhizobiales</i>	TAS [70, 71]
		Family <i>Rhizobiaceae</i>	TAS [72, 73]
		Genus <i>Ensifer</i>	TAS [74]
		Species <i>Ensifer meliloti</i>	[21]
		Strain: Mlalz-1 (= USDA 1984)	IDA
	Gram stain	Negative	IDA
	Cell shape	Rod	IDA
	Motility	Motile	IDA
	Sporulation	Non-sporulating	NAS
	Temperature range	10–40 °C	IDA
	Optimum temperature	25–30 °C	IDA
	pH range; Optimum	5–9.5; 6.5–8	IDA
	Carbon source	Varied	IDA
MIGS-6	Habitat	Soil; root nodule on host <i>Medicago laciniata</i> (L.) Miller	IDA
MIGS-6.3	Salinity	Tolerates 0 to 1% (w/v) % NaCl	TAS
MIGS-22	Oxygen requirement	Aerobic	IDA
MIGS-15	Biotic relationship	Free living, symbiotic	IDA
MIGS-14	Pathogenicity	Biosafety level 1	TAS [75]
MIGS-4	Geographic location	Guatiza, Lanzarote, Canary Islands, Spain	IDA
MIGS-5	Sample collection date	2007	IDA
MIGS-4.1	Latitude	29.074324	IDA
MIGS-4.2	Longitude	–13.479696	IDA
MIGS-4.3	Depth	5–10 cm	IDA
MIGS-4.4	Altitude	102 m	IDA

^aEvidence codes – IDA Inferred from Direct Assay, TAS Traceable Author Statement (i.e., a direct report exists in the literature), NAS Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). Evidence codes are from the Gene Ontology project [76, 77]

12611^T and *Ensifer terangae* LMG 7834^T [23] were only 97.3% similar.

Genome sequencing information

Genome project history

E. meliloti Mlalz-1 was selected for sequencing at the U.S. Department of Energy funded Joint Genome Institute as part of the GEBA-RNB project [17, 18]. The root nodule bacteria in this project were selected based on environmental and agricultural relevance to issues in global carbon cycling, alternative energy production, and biogeochemical importance. In particular, strain Mlalz-1 was chosen since it has strict host specificity for *M. laciniata*, which is suited for cultivation in arid environments [11]. The *E. meliloti* Mlalz-1 genome project is

deposited in the Genomes Online Database [24] and a high-quality permanent draft genome sequence (IMG Genome ID 2513237143) is deposited in IMG [25]. Sequencing, finishing and annotation were performed by the JGI. A summary of the project information is shown in Table 2.

Growth conditions and genomic DNA preparation

E. meliloti Mlalz-1 (= USDA 1984) was cultured on MAG solid media [26] for three days at 28 °C to obtain well grown, well separated colonies, then a single colony was selected from the plate and inoculated into 5 ml MAG broth media. The culture was grown for 48 h on a gyratory shaker (200 rpm) at 28 °C. Subsequently 1 ml was used to inoculate 50 ml of MAG and the cells were




Fig. 2 Comparison of the 16S rRNA gene sequences of *Ensifer meliloti* Mlalz-1 (shown in bold) and other *Ensifer* spp. type strains, based on aligned 16S rRNA gene sequences of 1389 bp. Of the 1389 sites, 1279 were constant and 48 were informative. There were eight gaps overall when *Mesorhizobium ciceri* bv *biserrulae* WSM1271 was included in the analysis. Phylogenetic analysis was done using MEGA, version 6.0 [61] after manually assembling the alignment by using GeneDoc version 2.6.001 [62]. *M. ciceri* bv *biserrulae* WSM1271 was used as an outgroup and the tree was assembled using the UPGMA algorithm based on the number of nucleotide differences. This approach was used since the potential for genetic recombination among the different 16S rRNA genes as reported by van Berkum [63] cannot be ignored. Bootstrap analysis [64] with 2000 permutations of the data set was done to assess support for the branch points. Strains with a genome sequencing project registered in GOLD [24] are *Ensifer adhaerens* Casida A^T, *M. ciceri* bv. *biserrulae* WSM1271 and Mlalz-1 and the GOLD ID is provided in place of the GenBank accession number

Table 2 Genome sequencing project information for *Ensifer meliloti* Mlalz-1

MIGS ID	Property	Term
MIGS-31	Finishing quality	High-quality permanent draft
MIGS-28	Libraries used	Illumina Standard PE
MIGS-29	Sequencing platforms	Illumina HiSeq 2000
MIGS-31.2	Fold coverage	748x
MIGS-30	Assemblers	Velvet version 1.1.04; ALLPATHS v. r39750
MIGS-32	Gene calling methods	Prodigal 1.4
	Locus Tag	A3CA [78]
	GenBank ID	ATZD00000000
	Genbank Date of Release	January 30 2012
	GOLD ID	Gp0010229 [79]
	BIOPROJECT	165,343
MIGS-13	Source Material Identifier	Mlalz-1 (=USDA 1984)
	Project relevance	Symbiotic N ₂ fixation, agriculture

incubated on a gyratory shaker (200 rpm) at 28 °C until an OD_{600nm} of 0.6 was reached. DNA was isolated from 50 ml of cells by Peter van Berkum according to the method described by van Berkum [26]. The final concentration of the DNA was set to 0.5 mg ml⁻¹.

Genome sequencing and assembly

The draft genome of *E. meliloti* Mlalz-1 was generated at the DOE Joint genome Institute (JGI) using Illumina technology [27]. An Illumina standard PE library was constructed and sequenced using the Illumina HiSeq 2000 platform that generated 35,720,836 reads totalling 4983 Mbp. All general aspects of library construction and sequencing were done at the JGI and details can be found on the JGI website [28]. All raw Illumina sequence data was passed through DUK, a filtering program developed at JGI, which removes known Illumina sequencing and library preparation artefacts (Mingkun L, Copeland A, Han J; unpublished). The following steps for assembly were: (1) filtered Illumina reads were assembled using Velvet (version 1.1.04) [29]; (2) 1–3 Kbp simulated paired end reads were created from Velvet contigs using wgsim (version 0.3.0) [30]; (3) Illumina reads were

assembled with simulated read pairs using Allpaths-LG (version r39750) [31]. Parameters for the assembly steps were 1) Velvet: `-v -s 51 -e 71 -i 2 -t 1 -f "-shortPaired -fastq $FASTQ" -o "-ins_length 250 -min_contig_lgth 500"` for Velvet and 2) wgsim: `-e 0-1 76-2 76 -r 0 -R 0 -X 0`. The final draft assembly contained 100 contigs in 99 scaffolds. The total size of the genome is 6.7 Mbp and the final assembly is based on 4983 Mbp of Illumina data, which provides an average of 748× coverage of the genome.

Genome annotation

Genes were identified using Prodigal [32], as part of the DOE-JGI genome annotation pipeline [33, 34]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information non-redundant database, UniProt, TIGRFam, Pfam, KEGG, COG, and InterPro databases. The tRNAScanSE tool [35] was used to find tRNA genes, whereas ribosomal RNA genes were found by searches against models of the ribosomal RNA genes built from SILVA [36]. Other non-coding RNAs such as the RNA components of the protein secretion complex and the RNase P were identified by searching the genome for the corresponding Rfam profiles using INFERNAL [37]. Additional gene prediction analysis and manual functional annotation was done within the Integrated Microbial Genomes-Expert Review platform [38] developed by the Joint Genome Institute, Walnut Creek, CA, USA.

Genome properties

The genome is 6,664,116 bp with 62.16% GC content (Table 3) and comprised of 99 scaffolds. From a total of 6388 genes, 6314 were protein encoding and 74 RNA

Table 3 Genome statistics for *Ensifer meliloti* Mlalz-1

Attribute	Value	% of Total
Genome size (bp)	6,664,116	100.00
DNA coding (bp)	5,754,332	86.35
DNA G + C (bp)	4,142,407	62.16
DNA scaffolds	99	100.00
Total genes	6388	100.00
Protein-coding genes	6314	98.84
RNA genes	74	1.16
Pseudo genes	0	0.00
Genes in internal clusters	1054	16.50
Genes with function prediction	5080	79.52
Genes assigned to COGs	4659	72.93
Genes with Pfam domains	5317	83.23
Genes with signal peptides	555	8.69
Genes with transmembrane helices	1440	22.54
CRISPR repeats	0	0.00

only encoding genes. Most genes (79.52%) were assigned a putative function whilst the remaining genes were annotated as hypothetical. The distribution of genes into COGs functional categories is presented in Table 4.

Insights from the genome sequence

E. meliloti Mlalz-1 is one of seven strains of *E. meliloti* that have been sequenced from the GEBA-RNB genome sequencing projects [17]. On the basis of 16S rRNA sequence identity, strain Mlalz-1 is closely related to *E. meliloti* IAM 12611^T (= LMG 6133^T), *E. medicae* A 321^T (= LMG 19920^T) and *E. numidicus* ORS 1407^T. As the genomes of these type strains have not been sequenced or are not publically available, gANI values [39]

Table 4 Number of genes of *Ensifer meliloti* Mlalz-1 associated with the general COG functional categories

Code	Value	%age	Description
J	217	4.09	Translation, ribosomal structure and biogenesis
A	0	0.00	RNA processing and modification
K	466	8.77	Transcription
L	122	2.3	Replication, recombination and repair
B	1	0.02	Chromatin structure and dynamics
D	39	0.73	Cell cycle control, cell division, chromosome partitioning
Y	0	0.00	Nuclear structure
V	117	2.20	Defense mechanisms
T	216	4.07	Signal transduction mechanisms
M	301	5.67	Cell wall/membrane/envelope biogenesis
N	72	1.36	Cell motility
Z	0	0.00	Cytoskeleton
W	33	0.62	Extracellular structures
U	74	1.39	Intracellular trafficking, secretion, and vesicular transport
O	206	3.88	Posttranslational modification, protein turnover, chaperones
C	358	6.74	Energy production and conversion
G	555	10.45	Carbohydrate transport and metabolism
E	584	10.99	Amino acid transport and metabolism
F	116	2.18	Nucleotide transport and metabolism
H	242	4.56	Coenzyme transport and metabolism
I	220	4.14	Lipid transport and metabolism
P	279	5.25	Inorganic ion transport and metabolism
Q	159	2.99	Secondary metabolite biosynthesis, transport and catabolism
R	551	10.37	General function prediction only
S	348	6.55	Function unknown
X	36	0.68	Mobilome: prophages, transposons
-	1729	27.07	Not in COGS

had to be compared with other fully sequenced *Ensifer* strains (Table 5). *E. meliloti* Mlalz-1 currently forms a gANI clique with other *E. meliloti* strains (gANI values $\geq 98.14\%$), compared with gANI values of $\leq 87.9\%$ with the finished genomes of other *Ensifer* strains. This supports the classification of strain Mlalz-1 as an *E. meliloti* strain, in accordance with the defined species affiliation cut-off value of 96.5% gANI [39]. The total genome size of strain Mlalz-1 is 6.6 Mbp, which falls within the expected size range of 6.6–8.9 Mbp for *E. meliloti*. The genome architecture of *E. meliloti* consists of a chromosome and the two symbiotic megaplastids pSymA and pSymB [20]. Replication of a plasmid is initiated by the replication protein encoded by *repC*, which is present as a single copy on *E. meliloti* pSymA and pSymB. The *E. meliloti* Mlalz-1 genome carried 2 *repC* loci (A3CADRAFT_00120 and A3CADRAFT_01676) with highest encoded protein identity to RepC proteins of *E. meliloti* strains. Mlalz-1 A3CADRAFT_00120 RepC1 had highest identity (98.10%) to the RepC1 protein encoded by SMB20044 on pSymB of *E. meliloti* 1021. *E. meliloti* Mlalz-1 A3CADRAFT_01676 RepC2 had highest identity (99.00%) to the RepC2 protein encoded by SMA2391 on pSymA of *E. meliloti* 1021. This indicated the presence of two megaplastids in strain Mlalz-1, and revealed that strain Mlalz-1 has a similar genome architecture to that of *E. meliloti* 1021.

Extended insights

All 29 *E. meliloti* strains within the gANI clique share a core set of 4948 orthologous genes, using cut off values of $1e-5$ and 30% minimum protein identity. *E. meliloti* Mlalz-1 contains 176 unique genes, 96 (54.5%) of which encode hypothetical proteins. The unique genes include those encoding the components of a T2SS, located on scaffold A3CADRAFT_scaffold_5.6 (Fig. 3a), as well as genes that encode a DNA methyltransferase and a NitT/TauT family transport system. These T2SS components

form part of a unique COG profile generated for Mlalz-1 (Table 6). The T2SS secretion system is used to translocate a wide range of proteins from the periplasm across the outer membrane [40]. Although T2SS genes are not found in other *E. meliloti* strains or in the *Ensifer fredii* strains GR64 and USDA 257, they are present in the genomes of the *E. fredii* strains HH103 and NGR234, in a similar gene arrangement to that observed in *E. meliloti* Mlalz-1 [41, 42] (Fig. 3b). Generally, the T2SS gene cluster is comprised of 12–15 genes, and strain Mlalz-1 contains the 12 required genes *gspDOGLMCKEFHII* necessary for a functional T2SS, but lacks the *gspS* gene found only in certain genera [43] (Fig. 3c).

In common with some other *E. meliloti* strains, strain Mlalz-1 contains several genes encoding phage components. The PHASTER algorithm [44] was used to identify two resident prophages, present on scaffold A3CADRAFT_scaffold_4.5: one that was incomplete (Prophage Region 1) and one that was intact (Prophage Region 2) (Fig. 4). The proteins encoded by Prophage Region 1 (11.4 kb) and Prophage Region 2 (55 kb) were most closely related to the phage proteins of PHAGE_Mycoba_Catalina_NC031238 and PHAGE_Sinorh_phiLM21_NC_029046, respectively.

The Mlalz-1 genome also contains acid-tolerance or acid-responsive genes that are orthologous to the genes identified in the comparatively acid tolerant strain *E. medicae* WSM419. Acid-tolerance or acid-responsive genes identified in Mlalz-1 include *actA* (*Int*), *actP*, *actR*, *actS*, *phrR*, *exoR*, *exoH*, *lpiA*, *acvB*, *degP1*, *mdh3*, *fbaB*, *groS*, *kdpB*, *kdpC*, *fixN2* and *fixO2* [45–52] (Additional file 2: Table S2). It is notable that strain Mlalz-1 is unique among the sequenced *Ensifer* strains since it contains two versions of the highly acid-induced *lpiA-acvB* operon. One operon (A3CADRAFT_01189–A3CADRAFT_01190) is found on scaffold A3CADRAFT_scaffold_3.4, in a gene region that is conserved in other *E. meliloti* (sequence similarity >98%) and is located on the

Table 5 Pairwise gANI comparisons of selected finished genomes of sequenced *Ensifer* strains

Strain	Gold ID: Gp	Casida A	USDA 257	WSM 419	1021	AK83	BL225C	GR4	Mlalz-1	Rm41	SM11
<i>E. adhaerens</i> Casida A	0094824	100	80.5	79.06	80.12	80.11	80.06	80.01	80.08	80.03	80.06
<i>E. fredii</i> USDA 257	0005169	80.5	100	81.89	83.26	83.24	83.25	83.20	83.14	83.33	83.22
<i>E. medicae</i> WSM419	0000117	79.06	81.93	100	88.18	88.13	88.26	88.24	87.90	88.14	88.26
<i>E. meliloti</i> 1021	0000726	80.12	83.26	88.19	100	99.36	99.62	99.41	98.80	99.24	99.43
<i>E. meliloti</i> AK83	0006695	80.08	83.25	88.16	99.36	100	99.33	99.14	98.60	99.38	99.33
<i>E. meliloti</i> BL225C	0006560	80.06	83.25	88.28	99.62	99.33	100	99.44	98.81	99.26	99.39
<i>E. meliloti</i> GR4	0020501	80.01	83.23	88.26	99.41	99.14	99.43	100	98.81	99.05	99.25
<i>E. meliloti</i> Mlalz-1	0010229	80.11	83.15	87.91	98.80	98.59	99.81	98.81	100	98.59	98.66
<i>E. meliloti</i> Rm41	0025853	80.05	83.36	88.11	99.26	99.39	99.25	99.06	98.59	100	99.33
<i>E. meliloti</i> SM11	0006018	80.05	83.23	88.29	99.45	99.33	99.39	99.26	98.67	99.32	100

For *E. meliloti* Mlalz-1, gANI values above the microbial species delineation cutoff value of 96.5% [39] are in bold font




Fig. 3 **a** Map of *Ensifer meliloti* Mlalz-1: A3CADRAFT_scaffold_5.6. **b** Genetic organization of the T2SS clusters in *E. meliloti* Mlalz-1 (*Emel* Mlalz-1), *Ensifer fredii* HH103 (*Efre* HH103) and *Ensifer fredii* NGR234 (*Efre* NGR234). J, *gspJ*; I, *gspI*; H, *gspH*; F, *gspF*; E, *gspE*; K, *gspK*; C, *gspC*; M, *gspM*; L, *gspL*; G, *gspG*; O, *gspO*; D, *gspD*. **c** Schematics of the T2SS of Gram-negative bacteria [43]. The secretin, GspD (A3CADRAFT_01569); the polytopic protein, GspF (A3CADRAFT_01561); the cytoplasmic ATPase, GspE (A3CADRAFT_01562); the major pseudopilin component, GspG (A3CADRAFT_01567); the minor pseudopilins, GspH (A3CADRAFT_01560), GspI (A3CADRAFT_01559), GspJ (A3CADRAFT_01558) and GspK (A3CADRAFT_01563); the bitopic proteins, GspL (A3CADRAFT_01566), GspC (A3CADRAFT_01564) and GspM (A3CADRAFT_01565); the peptidase, GspO (A3CADRAFT_01568) (GspS is absent from Mlalz-1); OM, outer membrane; PG, peptidoglycan; IM, inner membrane

chromosome of the fully sequenced *E. meliloti* 1021. The second version of the *lpiA-acvB* operon (A3CADRAFT_05694-A3CADRAFT_05695) is located on A3CADRAFT_scaffold_47.48, in a gene region that is conserved in *E. medicae* genomes (sequence similarity >96%) and is located on the pSMED02 symbiotic plasmid of the fully sequenced *E. medicae* WSM419. The regulatory gene *fsrR*, required for the acid activated expression of *lpiA* in *E. medicae* WSM419 [53], is located upstream of A3CADRAFT_05694 in strain Mlalz-1. This regulatory gene is absent from the

A3CADRAFT_01190 gene region, and from the *lpiA-acvB* gene regions of all other *E. meliloti* sequenced genomes. These findings suggest that *E. meliloti* Mlalz-1 acquired the plasmid-borne *lpiA-acvB* operon and associated *fsrR* regulatory gene by lateral transfer from an *E. medicae* strain.

Essential symbiotic (*nod*, *nif* and *fix*) genes identified in the *E. meliloti* Mlalz-1 genome (Additional file 2: Table S3 and S4) are located in several clusters on the following scaffolds: A3CADRAFT_scaffold_54.55 (Fig. 5a), A3CADRAFT_scaffold_61.62 (Fig. 5b), A3CADRAFT_scaffold_63.64 (Fig. 5c), A3CADRAFT_scaffold_71.72 (Fig. 5d)

Table 6 List of the unique COGs in *Ensifer meliloti* Mlalz-1


COG	Name	Locus Tag	Gene symbol	Protein function
0393	Uncharacterized conserved protein YbjQ, UPF0145 family	A3CADRAFT_01446		Unknown
4970	Tfp pilus assembly protein FimT	A3CADRAFT_01560	<i>gspH</i>	T2SS
1459	Type II secretory pathway, component PulF	A3CADRAFT_01561	<i>gspF</i>	T2SS
2804	Type II secretory pathway ATPase GspE/PulE or T4P pilus assembly pathway ATPase PilB	A3CADRAFT_01562	<i>gspE</i>	T2SS
3156	Type II secretory pathway, component PulK	A3CADRAFT_01563	<i>gspK</i>	T2SS
3166	Tfp pilus assembly protein PilN	A3CADRAFT_01566	<i>gspL</i>	T2SS
2165	Type II secretory pathway, pseudopilin PulG	A3CADRAFT_01567	<i>gspG</i>	T2SS
1450	Type II secretory pathway component GspD/PulD (secretin)	A3CADRAFT_01569	<i>gspD</i>	T2SS
2189	Adenine specific DNA methylase Mod	A3CADRAFT_02454	<i>yhdJ</i>	DNA methyltransferase
4705	Uncharacterized membrane-anchored protein	A3CADRAFT_05679		Membrane protein
4089	Uncharacterized membrane protein	A3CADRAFT_05685		Membrane protein
2021	Homoserine acetyltransferase	A3CADRAFT_06155		Homoserine acetyltransferase

and A3CADRAFT_scaffold_74.75 (Fig. 5e). Nodulation of *M. laciniata* has been shown to require a specific *nodC* allele [14]. The *nodC* gene of strain Mlalz-1 has highest sequence identity ($\geq 98\%$) with *nodC* of other *M. laciniata*-nodulating *Ensifer* strains in the NCBI database, whereas there is a lower sequence identity ($\leq 93\%$) with *nodC* of *Ensifer* strains that nodulate other *Medicago* species. Nodulation of *Medicago* hosts requires Nod factors that are sulfated at the reducing terminus and acylated at the non-reducing terminus, with a polyunsaturated fatty acyl tail [54, 55]. The NodH sulfotransferase, together with the NodP and NodQ sulfate-activating complex, are required for Nod factor sulfation [56, 57]. Activity of NodL results in O-acetylation of the Nod factor [58], while NodE and NodF produce the specific polyunsaturated fatty acyl tail [55, 59].

Strain Mlalz-1 would appear to be typical of *Ensifer* strains that nodulate *Medicago* species since the *nodeE*, *nodL* and *nodHPQ* genes that are required for these specific decorations of the Nod factor are present in the genome. *E. meliloti* Mlalz-1 also possesses the three *nodD* genes that mediate host-specific activation of *nodABC* in the symbiotic interactions of *E. meliloti* with *Medicago* [60].

Conclusions

E. meliloti Mlalz-1 is a rhizobial strain that is able to nodulate and fix nitrogen with the highly specific host *M. laciniata*. Although the 16S rRNA gene sequence divergence was insufficient to differentiate strain Mlalz-1 from *E. meliloti*, *E. medicae* or *E. numidicus*, a gANI value of 98.8% with the genome of *E. meliloti* 1021,








a Emel Mlalz-1: A3CADRAFT_scaffold_54.55**b** Emel Mlalz-1: A3CADRAFT_scaffold_61.62**c** Emel Mlalz-1: A3CADRAFT_scaffold_63.64**d** Emel Mlalz-1: A3CADRAFT_scaffold_71.72**e** Emel Mlalz-1: A3CADRAFT_scaffold_74.75

Fig. 5 Graphical map of the scaffolds; **a** A3CADRAFT_scaffold_54.55, **b** A3CADRAFT_scaffold_61.62, **(c)** A3CADRAFT_scaffold_63.64, **d** A3CADRAFT_scaffold_71.72 and **e** A3CADRAFT_scaffold_74.75 of *Ensifer meliloti* Mlalz-1 showing the location of common nodulation (*nod*) and fixation (*nif* and *fix*) genes within the symbiotic regions of this strain. From bottom to the top of the scaffold map: Genes on reverse strand (color by COG categories as denoted by the IMG platform), genes on forward strand (color by COG categories), RNA genes (tRNAs green, sRNAs red, other RNAs black), GC content, GC skew

compared with 87.9% with the genome of *E. medicae* WSM419 identifies strain Mlalz-1 as *E. meliloti*. Nodulation of *M. laciniata* has been shown to be dependent on the presence of a specific *nodC* allele, which also is present in the genome of *E. meliloti* Mlalz-1, based on a 98% sequence identity with the *nodC* of other *M. laciniata*-nodulating *Ensifer* strains [14]. However, strain Mlalz-1 is unique among sequenced *E. meliloti* strains in possessing genes encoding components of a T2SS and in having two versions of the adaptive acid tolerance response *lpiA-acvB* operon. The second copy of the *E. meliloti* Mlalz-1 *lpiA-acvB* operon has highest sequence identity (>96%) with that of sequenced *E. medicae* strains, which infers horizontal gene transfer of this region from *E. medicae*.

Additional files

Additional file 1: Table S1. Associated MIGS record for *Ensifer meliloti* Mlalz-1. (DOCX 52 kb)

Additional file 2: Table S2–S4. Table S2. Acid responsive gene orthologs present in *Ensifer* strains. **Table S3.** The nodulation genes of *Ensifer meliloti* Mlalz-1. **Table S4.** The nitrogen fixation genes of *Ensifer meliloti* Mlalz-1. (DOCX 65 kb)

Abbreviations

½LA: Half strength Lupin Agar; gANI: Genome-wide average nucleotide identity; GEBA-RNB: Genomic Encyclopedia for Bacteria and Archaea-Root Nodule Bacteria; IMG: Integrated Microbial Genomes; T2SS: Type II Secretion System; TY: Tryptone-yeast extract

Acknowledgements

We thank Gordon Thompson (Murdoch University) for the preparation of SEM and TEM photos. MLB thanks Alfredo Reyes-Betancort, from the Orotava Botanical Garden (Tenerife), for providing *M. laciniata* seeds.

Funding

This work was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under contract No. DE-AC02-05CH11231. We gratefully acknowledge the funding received from the Curtin University Sustainability Policy Institute and from Murdoch University's Small Research Grants Scheme in 2016.

Authors' contributions

MLB and EV isolated the strain and provided project metadata. PvB supplied the strain, the DNA and the background information for this project and participated in drafting the manuscript. PE curated the strain and performed sequence analysis of 16S rRNA and *nodC* genes. RT supplied DNA to JGI and performed all imaging. JA provided symbiotic phenotype data. WAMO, JA and WR performed bioinformatics analyses and drafted the paper, MB and NB provided financial support, and MG, RS, TBKR, NI, TW, AP, VM and NK were involved in sequencing the genome and/or editing the final paper. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹School of Veterinary and Life Sciences, Murdoch University, Murdoch, WA, Australia. ²U.S. Department of Agriculture, Soybean Genomics and

Improvement Laboratory, Beltsville Agricultural Research Center, 10300 Baltimore Avenue, Bldg. 006, Beltsville, MD 20705, USA. ³Departamento de Bioquímica, Microbiología, Biología Celular y Genética, Universidad de La Laguna, Tenerife, Spain. ⁴Departamento de Microbiología y Genética and Instituto Hispanoluso de Investigaciones Agrarias (CIALE), Universidad de Salamanca, Salamanca, Spain. ⁵Curtin University Sustainability Policy Institute, Curtin University, Bentley, WA, Australia. ⁶DOE Joint Genome Institute, Walnut Creek, CA, USA. ⁷Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. ⁸Department of Biology, Faculty of Science, University of Jeddah, Jeddah, Saudi Arabia.

Received: 30 March 2017 Accepted: 14 September 2017

Published online: 25 September 2017

References

- Carlsson G, Huss-Danell K. Nitrogen fixation in perennial forage legumes in the field. *Plant Soil*. 2003;253:353–72.
- Peoples MB, Brockwell J, Hunt JR, Swan AD, Watson L, Hayes RC, Li GD, Hackney B, Nuttall JG, Davies SL, Fillery IRP. Factors affecting the potential contributions of N₂ fixation by legumes in Australian pasture systems. *Crop Pasture Sci*. 2012;63:759–86.
- Unkovich MJ, Baldock J, Peoples MB. Prospects and problems of simple linear models for estimating symbiotic N₂ fixation by crop and pasture legumes. *Plant Soil*. 2010;329:75–89.
- UNDP 2015 [http://www.undp.org/content/undp/en/home/sdoverview/post-2015-development-agenda.html]. Accessed 22 Sept 2017.
- Béna G, Lyet A, Huguet T, Olivieri I. *Medicago*–*Sinorhizobium* symbiotic specificity evolution and the geographic expansion of *Medicago*. *J Evol Biol*. 2005;18:1547–58.
- Rome S, Fernandez MP, Brunel B, Normand P, Cleyet-Marel JC. *Sinorhizobium medicae* sp. nov., isolated from annual *Medicago* spp. *Int J Syst Bacteriol*. 1996;46:972–80.
- Jones KM, Kobayashi H, Davies BW, Taga ME, Walker GC. How rhizobial symbionts invade plants: the *Sinorhizobium*–*Medicago* model. *Nature Rev Microbiol*. 2007;5:619–33.
- Jordan DC. Reduction of the nodulation barrier in *Medicago laciniata* by alteration of the root temperature. *Plant Soil*. 1981;61:93–111.
- Small E. *Alfalfa and relatives: Evolution and classification of Medicago*. Ottawa: NRC Research Press; 2010.
- Young RR, Croft PH, Sandral GA. Variation in flowering times and agronomic characteristics of *Medicago laciniata* (L.) miller collected from diverse locations in new South Wales. *Aust J Exp Agric*. 1992;32:59–63.
- Yousfi N, Sihem N, Ramzi A, Abdelly C. Growth, photosynthesis and water relations as affected by different drought regimes and subsequent recovery in *Medicago laciniata* (L.) populations. *J Plant Biol*. 2016;59:33–43.
- Brockwell J, Hely FW. Symbiotic characteristics of *Rhizobium meliloti*: an appraisal of the systematic treatment of nodulation and nitrogen fixation interactions between hosts and rhizobia of diverse origins. *Aust J Agr Econ*. 1966;17:885–9.
- Villegas MDC, Rome S, Mauré L, Domergue O, Gardan L, Bailly X, Cleyet-Marel J-C, Brunel B. Nitrogen-fixing sinorhizobia with *Medicago laciniata* constitute a novel biovar (bv. *Medicaginis*) of *S. meliloti*. *Syst Appl Microbiol*. 2006;29:526–38.
- Barran LR, Bromfield ES, Brown DC. Identification and cloning of the bacterial nodulation specificity gene in the *Sinorhizobium meliloti*–*Medicago laciniata* symbiosis. *Can J Microbiol*. 2002;48:765–71.
- van Berkum P, Badri Y, Elia P, Aouani ME, Eardly BD. Chromosomal and symbiotic relationships of rhizobia nodulating *Medicago truncatula* and *M. laciniata*. *Appl Environ Microbiol*. 2007;73:7597–604.
- Mnasri B, Badri Y, Saïdi S, de Lajudie P, Mhamdi R. Symbiotic diversity of *Ensifer meliloti* strains recovered from various legume species in Tunisia. *Syst Appl Microbiol*. 2009;32:583–92.
- Reeve W, Ardley J, Tian R, Eshragi L, Yoon JW, Ngamwisetkun P, Seshadri R, Ivanova NN, Kyrpidis NC. A genomic encyclopedia of the root nodule bacteria: assessing genetic diversity through a systematic biogeographic survey. *Stand Genomic Sci*. 2015;10:14.
- Seshadri R, Reeve WG, Ardley JK, Tennessen K, Woyke T, Kyrpidis NC, Ivanova NN. Discovery of novel plant interaction determinants from the genomes of 163 root nodule bacteria. *Sci Rep*. 2015;5:16825.
- Howieson JG, Dilworth MJ, editors. *Working with Rhizobia*. Canberra, Australia: Australian Centre for International Agricultural Research (ACIAR); 2016.

20. Bailly X, Olivieri I, Brunel B, Cleyet-Marel JC, Béna G. Horizontal gene transfer and homologous recombination drive the evolution of the nitrogen-fixing symbionts of *Medicago* species. *J Bacteriol.* 2007;189:5223–36.
21. Delajudie P, Willems A, Pot B, Dewettinck D, Maestrojuan G, Neyra M, Collins MD, Dreyfus B, Kersters K, Gillis M. Polyphasic taxonomy of rhizobia: emendation of the genus *Sinorhizobium* and description of *Sinorhizobium meliloti* comb. nov., *Sinorhizobium saheli* sp. nov., and *Sinorhizobium teranga* sp. nov. *Int J Syst Bacteriol.* 1994;44:715–33.
22. Young JM. The genus name *Ensifer* Casida 1982 takes priority over *Sinorhizobium* Chen et al. 1988, and *Sinorhizobium morelense* Wang et al. 2002 is a later synonym of *Ensifer adhaerens* Casida 1982. Is the combination "*Sinorhizobium adhaerens*" (Casida 1982) Willems et al. 2003 legitimate? Request for an opinion. *Int J Syst Evol Microbiol.* 2003;53:2107–10.
23. Merabet C, Martens M, Mahdhi M, Zakhia F, Sy A, Le Roux C, Domergue O, Coopman R, Bekki A, Mars M, et al. Multilocus sequence analysis of root nodule isolates from *Lotus arabicus* (Senegal), *Lotus creticus*, *Argyrolobium uniflorum* and *Medicago sativa* (Tunisia) and description of *Ensifer numidicus* sp. nov. and *Ensifer garamanticus* sp. nov. *Int J Syst Evol Microbiol.* 2010;60:664–74.
24. Reddy TBK, Thomas AD, Stamatis D, Bertsch J, Isbandi M, Jansson J, Mallajosyula J, Pagani I, Lobos EA, Kyrpidis NC. The genomes OnLine database (GOLD) v.5: a metadata management system based on a four level (meta)genome project classification. *Nucleic Acids Res.* 2015;43:D1099–106.
25. Markowitz VM, Chen IM, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A, Huang J, Woyke T, Huntemann M, et al. IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res.* 2014;42:D560–7.
26. van Berkum P. Evidence for a third uptake hydrogenase phenotype among the soybean bradyrhizobia. *Appl Environ Microbiol.* 1990;56:3835–41.
27. Bennett S. Solexa Ltd. Pharmacogenomics. 2004;5:433–8.
28. Joint Genome Institute website [<http://jgi.doe.gov/>]. Accessed 22 Sept 2017.
29. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 2008;18:821–9.
30. GitHub - lh3/wgsm: Reads simulator [<https://github.com/lh3/wgsm>]. Accessed 22 Sept 2017.
31. Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, et al. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A.* 2011;108:1513–8.
32. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics.* 2010;11:119.
33. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Páez-Espino D, Palaniappan K, Szeto E, Pillay M, Chen IM, Pati A, et al. The standard operating procedure of the DOE-JGI microbial genome annotation pipeline (MGAP v.4). *Stand Genomic Sci.* 2015;10:86.
34. Chen IM, Markowitz VM, Chu K, Anderson I, Mavromatis K, Kyrpidis NC, Ivanova NN. Improving microbial genome annotations in an integrated database context. *PLoS One.* 2013;8:e54859.
35. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 1997;25:955–64.
36. Pruesse E, Quast C, Knittel K, Bdm F, Ludwig W, Peplies J, Glöckner FO. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res.* 2007;35:7188–96.
37. Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics.* 2013;29:2933–5.
38. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpidis NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics.* 2009;25:2271–8.
39. Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavromatis K, Kyrpidis NC, Pati A. Microbial species delineation using whole genome sequences. *Nucleic Acids Res.* 2015;43:6761–71.
40. Korotkov KV, Sandkvist M, Hol WG. The type II secretion system: biogenesis, molecular architecture and mechanism. *Nature Rev Microbiol.* 2012;10:336–51.
41. Kryciak D, Orbegoso MR, Schmeisser C, Streit WR. Molecular keys to broad host range in *Sinorhizobium fredii* NGR234, USDA257 and HH103. In: De Bruijn FJ, editor. *Biological nitrogen fixation*. Volume 2. Hoboken NJ: Wiley-Blackwell; 2015. p. 325–36.
42. Schmeisser C, Liesegang H, Kryciak D, Bakkou N, Le Quééré A, Wollherr A, Heinemeyer I, Morgenstern B, Pommerening-Röser A, Flores M, et al. *Rhizobium* sp. strain NGR234 possesses a remarkable number of secretion systems. *Appl Environ Microbiol.* 2009;75:4035–45.
43. Costa TR, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M, Waksman G. Secretion systems in gram-negative bacteria: structural and mechanistic insights. *Nature Rev Microbiol.* 2015;13:343–59.
44. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res.* 2016;44:W16–21.
45. Tiwari RP, Reeve WG, Dilworth MJ, Glenn AR. An essential role for *actA* in acid tolerance of *Rhizobium meliloti*. *Microbiology.* 1996;142:601–10.
46. Tiwari RP, Reeve WG, Dilworth MJ, Glenn AR. Acid tolerance in *Rhizobium meliloti* strain WSM419 involves a two-component sensor-regulator system. *Microbiology.* 1996;142:1693–704.
47. Tiwari RP, Reeve WG, Fenner BJ, Dilworth MJ, Glenn AR, Howieson JG. Probing for pH-regulated genes in *Sinorhizobium medicae* using transcriptional analysis. *J Mol Microbiol Biotechnol.* 2004;7:133–9.
48. Reeve WG, Dilworth MJ, Tiwari RP, Glenn AR. Regulation of exopolysaccharide production in *Rhizobium leguminosarum* biovar viciae WSM710 involves *exoR*. *Microbiology.* 1997;143:1951–8.
49. Reeve WG, Tiwari RP, Guerreiro N, Stubbs J, Dilworth MJ, Glenn AR, Rolfe BG, Djordjevic MA, Howieson JG. Probing for pH-regulated proteins in *Sinorhizobium medicae* using proteomic analysis. *J Mol Microbiol Biotechnol.* 2004;7:140–7.
50. Reeve WG, Tiwari RP, Kale NB, Dilworth MJ, Glenn AR. ActP controls copper homeostasis in *Rhizobium leguminosarum* bv. Viciae and *Sinorhizobium meliloti* preventing low pH-induced copper toxicity. *Mol Microbiol.* 2002;43:981–91.
51. Reeve WG, Tiwari RP, Wong CM, Dilworth MJ, Glenn AR. The transcriptional regulator gene *phrR* in *Sinorhizobium meliloti* WSM419 is regulated by low pH and other stresses. *Microbiology.* 1998;144:3335–42.
52. Glenn AR, Reeve WG, Tiwari RP, Dilworth MJ. Acid tolerance in root nodule bacteria. In: Chadwick DJ, Cardew G, editors. *Bacterial response to pH* Novartis Foundation symposium volume 221. 1999/04/20 edition. London: Wiley Publishing; 1999. p. 112–6.
53. Reeve WG, Brau L, Castelli J, Garau G, Sohlenkamp C, Geiger O, Dilworth MJ, Glenn AR, Howieson JG, Tiwari RP. The *Sinorhizobium medicae* WSM419 *lpiA* gene is transcriptionally activated by FsrR and required to enhance survival in lethal acid conditions. *Microbiology.* 2006;152:3049–59.
54. Lerouge P, Roche P, Faucher C, Maillet F, Truchet G, Promé JC, Dénarié J. Symbiotic host-specificity of *Rhizobium meliloti* is determined by a sulphated and acylated glucosamine oligosaccharide signal. *Nature.* 1990;344:781–4.
55. Dénarié J, Debelle F, Promé JC. *Rhizobium* lipo-chitoooligosaccharide nodulation factors: signaling molecules mediating recognition and morphogenesis. *Annu Rev Biochem.* 1996;65:503–35.
56. Schultze M, Staehelin C, Röhrig H, John M, Schmidt J, Kondorosi E, Schell J, Kondorosi A. In vitro sulfoltransferase activity of *Rhizobium meliloti* NodH protein: lipo-chitoooligosaccharide nodulation signals are sulfated after synthesis of the core structure. *Proc Natl Acad Sci U S A.* 1995;92:2706–9.
57. Schwedock JS, Liu C, Leyh TS, Long SR. *Rhizobium meliloti* NodP and NodQ form a multifunctional sulfate-activating complex requiring GTP for activity. *J Bacteriol.* 1994;176:7055–64.
58. Ardourel M, Lortet G, Maillet F, Roche P, Truchet G, Promé JC, Rosenberg C. In *Rhizobium meliloti*, the operon associated with the *nod* box n5 comprises *nodL*, *noeA* and *noeB*, three host-range genes specifically required for the nodulation of particular *Medicago* species. *Mol Microbiol.* 1995;17:687–99.
59. Demont N, Debelle F, Aurelle H, Dénarié J, Promé JC. Role of the *Rhizobium meliloti* *nodF* and *nodE* genes in the biosynthesis of lipo-oligosaccharidic nodulation factors. *J Biol Chem.* 1993;268:20134–42.
60. Honma MA, Asomaning M, Ausubel FM. *Rhizobium meliloti* *nodD* genes mediate host-specific activation of *nodABC*. *J Bacteriol.* 1990;172:901–11.
61. Tamura K, Stecher G, Peterson D, Filipiński A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol.* 2013;30:2725–9.
62. Nicholas KB, Nicholas HB, Deerfield DW. GeneDoc: analysis and visualization of genetic variation. *EMBN News.* 1997;4:14.
63. van Berkum P, Terefework Z, Paulin L, Suomalainen S, Lindström K, Eardly BD. Discordant phylogenies within the *rm* loci of rhizobia. *J Bacteriol.* 2003;185:2988–98.
64. Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution.* 1985;39:783–91.
65. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen M, Angiuoli SV, et al. Towards a richer description of our complete collection of genomes and metagenomes "minimum information about a genome sequence" (MIGS) specification. *Nature Biotechnol.* 2008;26:541–7.
66. Field D, Amaral-Zettler L, Cochrane G, Cole JR, Dawyndt P, Garrity GM, Gilbert J, Glöckner FO, Hirschman L, Karsch-Mizrachi I, et al. The genomic standards consortium. *PLoS Biol.* 2011;9:e1001088.

67. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, bacteria, and Eucarya. *Proc Natl Acad Sci U S A*. 1990;87:4576–9.
68. Garrity GM, Bell JA, Lilburn T. In: Garrity GM, Brenner DJ, Kreig NR, Staley JT, editors. Phylum XIV. Proteobacteria phyl. Nov. in *Bergey's manual of systematic bacteriology*. Volume 2. Second edition. New York: Springer - Verlag; 2005. p. 1.
69. Garrity GM, Bell JA, Lilburn T. In: Garrity GM, Brenner DJ, Kreig NR, Staley JT, editors. Class I. Alphaproteobacteria class. In *Bergey's manual of systematic bacteriology*. Second edition. New York: Springer - Verlag; 2005.
70. Euzéby J. Validation list no. 107. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol*. 2006;56:1–6.
71. Kuykendall LD. In: Garrity GM, Brenner DJ, Kreig NR, Staley JT, editors. Order VI. *Rhizobiales* ord. Nov. in *Bergey's manual of systematic bacteriology*. Second edition. New York: Springer - Verlag; 2005. p. 324.
72. Skerman VBD, McGowan V, Sneath PHA. Approved lists of bacterial names. *Int J Syst Bacteriol*. 1980;30:225–420.
73. Conn HJ. Taxonomic relationships of certain non-sporeforming rods in soil. *J Bacteriol*. 1938;36:320–1.
74. Casida LE. *Ensifer adhaerens* gen. Nov., sp. nov.: a bacterial predator of bacteria in soil. *Int J Syst Evol Microbiol*. 1982;32:339–45.
75. Biological Agents: Technical rules for biological agents [<http://www.baua.de/en/Topics-from-A-to-Z/Biological-Agents/TRBA/TRBA.html>]. Accessed 22 Sept 2017.
76. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene ontology: tool for the unification of biology. *The Gene Ontology Consortium Nature Genet*. 2000;25:25–9.
77. Guide to GO Evidence Codes [<http://geneontology.org/page/guide-go-evidence-codes>]. Accessed 22 Sept 2017.
78. Locus Tag [<https://www.ncbi.nlm.nih.gov/bioproject/?term=A3CA>]. Accessed 22 Sept 2017.
79. GOLD ID for *Ensifer meliloti* Mlalz-1 [<https://gold.jgi.doe.gov/projects?id=Gp0010229>]. Accessed 22 Sept 2017.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

